

ABSTRACT

W.J. Fokkens, V.J. Lund, J. Mullol et al., European Position Paper on Nasal Polyps 2007.
Rhinology 45; suppl. 20: 1-139.

* corresponding author: Wytse Fokkens, Department of Otorhinolaryngology, Amsterdam Medical Centre, PO box 22660, 1100 DD Amsterdam, The Netherlands. Email: w.j.fokkens@amc.nl

Rhinosinusitis is a significant and increasing health problem which results in a large financial burden on society. This evidence based position paper describes what is known about rhinosinusitis and nasal polyps, offers evidence based recommendations on diagnosis and treatment, and considers how we can make progress with research in this area.

Rhinitis and sinusitis usually coexist and are concurrent in most individuals; thus, the correct terminology is now rhinosinusitis. Rhinosinusitis (including nasal polyps) is defined as inflammation of the nose and the paranasal sinuses characterised by two or more symptoms, one of which should be either nasal blockage/obstruction/congestion or nasal discharge (anterior/posterior nasal drip), \pm facial pain/pressure, \pm reduction or loss of smell; and either endoscopic signs of polyps and/or mucopurulent discharge primarily from middle meatus and/or; oedema/mucosal obstruction primarily in middle meatus, and/or CT changes showing mucosal changes within the ostiomeatal complex and/or sinuses.

The paper gives different definitions for epidemiology, first line and second line treatment and for research.

Furthermore the paper describes the anatomy and (patho)physiology, epidemiology and predisposing factors, inflammatory mechanisms, evidence based diagnosis, medical and surgical treatment in acute and chronic rhinosinusitis and nasal polyposis in adults and children. Evidence based schemes for diagnosis and treatment are given for the first and second line clinicians. Moreover attention is given to complications and socio-economic cost of chronic rhinosinusitis and nasal polyps. Last but not least the relation to the lower airways is discussed.

Participants:

Wyske Fokkens, Chair. Department of Otorhinolaryngology, Amsterdam Medical Centre, Amsterdam

Valerie Lund, Co-Chair. Institute of Laryngology and Otolaryngology. University College London, London

Joaquim Mullol, Co-Chair. Rhinology Unit & Smell Clinic, ENT Department, Hospital Clínic – IDIBAPS, Barcelona, Spain

Claus Bachert. Upper Airway Research Laboratory, Department of Otorhinolaryngology, Ghent University, Belgium

Noam Cohen, Department of Otorhinolaryngology: Head and Neck Surgery, University of Pennsylvania, Philadelphia, USA

Roxanna Cobo, Department of Otolaryngology, Centro Médico Imbanaco, Cali, Colombia

Martin Desrosiers, Department of Otolaryngology-Head and Neck Surgery, University of Montreal, Montreal, Canada

Peter Hellings, Department of Otorhinolaryngology, University Hospitals Leuven, Catholic University Leuven, Leuven, Belgium

Mats Holmstrom, Department of Otorhinolaryngology, Uppsala University Hospital Uppsala, Sweden

Maija Hytönen, Department of Otorhinolaryngology, Helsinki University Central Hospital, Helsinki, Finland

Nick Jones, Department of Otorhinolaryngology, Head and Neck Surgery, Queen's Medical Centre, University of Nottingham, Nottingham, UK

Livije Kalogjera, Department of Otorhinolaryngology/Head and Neck Surgery, University Hospital "Sestre Milosrdnice", Zagreb, Croatia

David Kennedy, Department of Otorhinolaryngology: Head and Neck Surgery, University of Pennsylvania, Philadelphia, USA

Jean Michel Klossek, Department ENT and Head and neck surgery, CHU Poitiers: Univ Poitiers Hopital Jean Bernard, Poitiers, France

Marek Kowalski, Department of Immunology, Rheumatology and Allergy, Medical University of Lodz, Lodz, Poland

Eli Meltzer, Allergy and Asthma Medical Group and Research Center, University of California at San Diego, San Diego, California

Bob Naclerio, Section of Otolaryngology-Head and Neck Surgery, The Pritzker School of Medicine, The University of Chicago, Chicago, USA

Desiderio Passali, ENT Department, Policlinico Le Scotte - University of Siena, Siena, Italy

David Price, Dept of General Practice and Primary Care, University of Aberdeen, Aberdeen, UK

Herbert Riechelmann, University Hospital for Ear, Nose and Throat Diseases, University Hospital Center Ulm, Ulm, Germany

Glenis Scadding, Allergy & Medical Rhinology Department, Royal National TNE Hospital, London, UK.

Heinz Stammberger, University Ear, Nose and Throat Hospital, Graz, Austria

Mike Thomas, Department of General Practice and Primary Care, University of Aberdeen, Aberdeen, UK

Richard Voegels, Rhinology - Clinics, University of Sao Paulo Medical School, Sao Paulo, Brazil

De-Yun Wang. Department of Otolaryngology, National University of Singapore, Singapore

Acknowledgements:

The chairs of EP3OS would like to express their gratitude for the great help in preparing this document:

Fenna Ebbens, Amsterdam

Christos Georgalas, London

Hanneke de Bakker, Amsterdam

Josep Maria Guilemany, Barcelona

European Position Paper on Rhinosinusitis and Nasal Polyps 2007

CONTENTS

1	Introduction	5	8	Complications of rhinosinusitis and nasal polyps	80
			8-1	<i>Introduction</i>	80
			8-2	<i>Epidemiology of complications</i>	80
			8-3	<i>Orbital complications</i>	80
			8-4	<i>Endocranial complications</i>	01
			8-5	<i>Cavernous sinus thrombosis</i>	82
			8-6	<i>Osseous complications</i>	82
			8-7	<i>Unusual complications of rhinosinusitis</i>	83
2	Definition of rhinosinusitis and nasal polyps	6	9	Special considerations: Rhinosinusitis in children	84
2-1	<i>Introduction</i>	6	9-1	<i>Introduction</i>	84
2-2	<i>Clinical definition</i>	6	9-2	<i>Anatomy</i>	84
2-3	<i>Definition for use in epidemiology studies/ General Practice</i>	6	9-3	<i>Epidemiology and pathophysiology</i>	84
2-4	<i>Definition for research</i>	7	9-4	<i>Symptoms and signs</i>	85
3	Chronic rhinosinusitis with or without nasal polyps	8	9-5	<i>Clinical examination</i>	85
3-1	<i>Anatomy and (patho)physiology</i>	8	9-6	<i>Investigations</i>	85
3-2	<i>Rhinosinusitis</i>	8	9-7	<i>Management</i>	87
3-3	<i>Chronic rhinosinusitis with or without nasal polyps</i>	8	10	Chronic rhinosinusitis with or without nasal polyps in relation to the lower airways	90
4	Epidemiology and predisposing factors	10	10-1	<i>Introduction</i>	90
4-1	<i>Introduction.</i>	10	10-2	<i>Asthma and Chronic Rhinosinusitis without NP</i>	90
4-2	<i>Acute bacterial rhinosinusitis.</i>	10	10-3	<i>Asthma and Chronic Rhinosinusitis with NP</i>	91
4-3	<i>Factors associated with acute rhinosinusitis.</i>	10	10-4	<i>COPD and rhinosinusitis</i>	91
4-4	<i>Chronic rhinosinusitis (CRS) without nasal polyps</i>	12	11	Socio-economic cost of chronic rhinosinusitis and nasal polyps	92
4-5	<i>Factors associated with chronic rhinosinusitis (CRS) without nasal polyps</i>	12	11-1	<i>Direct Costs</i>	92
4-6	<i>Chronic rhinosinusitis with nasal polyps</i>	15	11-2	<i>Indirect Costs</i>	92
4-7	<i>Factors associated with Chronic rhinosinusitis with nasal polyps</i>	16	12	Outcomes measurements in research	94
4-8	<i>Epidemiology and predisposing factors for rhinosinusitis in children</i>	18	13	Evidence based schemes for diagnostic and treatment	95
4-9	<i>Conclusion</i>	18	13-1	<i>Introduction</i>	95
5	Inflammatory mechanisms in acute and chronic rhinosinusitis with or without nasal polyposis	19	13-2	<i>Introduction</i>	97
5-1	<i>Introduction</i>	19	13-3	<i>Evidence based management scheme for adults with acute rhinosinusitis</i>	98
5-2	<i>Acute rhinosinusitis</i>	19	13-4	<i>Evidence based management scheme for adults with chronic rhinosinusitis without nasal polyps</i>	100
5-3	<i>Chronic rhinosinusitis without nasal polyps</i>	20	13-5	<i>Evidence based schemes for therapy in children</i>	103
5-4	<i>Chronic rhinosinusitis with nasal polyps</i>	26	14	Research needs and priorities	105
5-5	<i>Aspirin sensitivity – Inflammatory mechanisms in acute and chronic rhinosinusitis</i>	32	14-1	<i>Epidemiology: Identifying the risk factors for development of CRS an NP</i>	105
5-6	<i>Conclusion</i>	34	14-2	<i>Beyond infection: New roles for bacteria</i>	105
6	Diagnosis	35	14-3	<i>Host response</i>	105
6-1	<i>Assessment of rhinosinusitis symptoms</i>	35	14-4	<i>Genetics</i>	105
6-2	<i>Examination</i>	37	14-5	<i>Clinical trials</i>	105
6-3	<i>Quality of Life</i>	40	15	GLOSARRY TERMS	107
7	Management	43	16	Information on QOL instruments:	108
7-1	<i>Treatment of rhinosinusitis with corticosteroids</i>	43	16-1	<i>General health status instruments:</i>	108
7-2	<i>Treatment of rhinosinusitis with antibiotics</i>	51	16-2	<i>Disease specific health status instruments:</i>	108
7-3	<i>Other medical management for rhinosinusitis</i>	56	17	Survey of published olfactory tests	109
7-4	<i>Evidence based surgery for rhinosinusitis</i>	64	18	References	111
7-5	<i>Influence of age concomitant diseases on sinus surgery outcome</i>	71			
7-6	<i>Complications of surgical treatment</i>	76			

1. Introduction

Rhinosinusitis is a significant health problem which seems to mirror the increasing frequency of allergic rhinitis and which results in a large financial burden on society ⁽¹⁻³⁾. Data on (chronic) rhinosinusitis are limited and the disease entity is badly defined. Therefore, the available data are difficult to interpret and extrapolate.

The last decade has seen the development of a number of guidelines, consensus documents and position papers on the epidemiology, diagnosis and treatment of rhinosinusitis and nasal polyposis ⁽⁴⁻⁷⁾. In 2005 the first European Position Paper on Rhinosinusitis and Nasal Polyps (EP3OS) was published ^(8,9). This first evidence based position paper was initiated by the European Academy of Allergology and Clinical Immunology (EAACI) to consider what was known about rhinosinusitis and nasal polyps, to offer evidence-based recommendations on diagnosis and treatment, and to consider how we can make progress with research in this area. The paper has been approved by the European Rhinologic Society.

Evidence-based medicine is an important method of preparing guidelines ^(10,11). Moreover, the implementation of guidelines is equally important.

Table 1-1. Category of evidence ⁽¹¹⁾


Ia	evidence from meta-analysis of randomised controlled trials
Ib	evidence from at least one randomised controlled trial
IIa	evidence from at least one controlled study without randomisation
IIb	evidence from at least one other type of quasi-experimental study
III	evidence from non-experimental descriptive studies, such as comparative studies, correlation studies, and case-control studies
IV	evidence from expert committee reports or opinions or clinical experience of respected authorities, or both

Table 1-2. Strength of recommendation

A	directly based on category I evidence
B	directly based on category II evidence or extrapolated recommendation from category I evidence
C	directly based on category III evidence or extrapolated recommendation from category I or II evidence
D	directly based on category IV evidence or extrapolated recommendation from category I, II or III evidence

Since the preparation of the first EP3OS document an increasing amount of evidence on the pathophysiology, diagnosis and treatment has been published (Figure 1).

Figure 1. Randomized controlled trials in chronic rhinosinusitis with or without nasal polyps. The number of trials in the last 5-6 years equals the number ever published before.



This revision is intended to be a state-of-the art review for the specialist as well as for the general practitioner:

- to update their knowledge of rhinosinusitis and nasal polyposis;
- to provide an evidence-based documented review of the diagnostic methods;
- to provide an evidence-based review of the available treatments;
- to propose a stepwise approach to the management of the disease;
- to propose guidance for definitions and outcome measurements in research in different settings.

In this revision new data have led to considerable increase in amount of available evidence and therefore to considerable changes in the schemes for diagnosis and treatment.

Moreover the whole document has been made more consistent, some chapters are significantly extended and others are added. Last but not least contributions from many other part of the world have attributed to our knowledge and understanding.

2. Definition of rhinosinusitis and nasal polyps

2-1 Introduction

Rhinitis and sinusitis usually coexist and are concurrent in most individuals; thus, the correct terminology is now rhinosinusitis. The diagnosis of rhinosinusitis is made by a wide variety of practitioners, including allergologists, otolaryngologists, pulmonologists, primary care physicians and many others. Therefore, an accurate, efficient, and accessible definition of rhinosinusitis is required. A number of groups have published reports on rhinosinusitis and its definition. In most of these reports definitions are based on symptomatology and duration of disease and a single definition is aimed at all practitioners^(4,5,12,13).

Due to the large differences in technical possibilities to diagnose and treat rhinosinusitis/nasal polyps by various disciplines, the need to differentiate between subgroups varies. On one hand the epidemiologist wants a workable definition that does not impose too many restrictions to study larger populations. On the other hand researchers in a clinical setting are in need of a set of clearly defined items that describes their patient population accurately and avoids the comparison of 'apples and oranges' in studies that relate to diagnosis and treatment. The taskforce tried to accommodate these different needs by offering definitions that can be applied in different circumstances. In this way the taskforce hopes to improve the comparability of studies, thereby enhancing the evidence based diagnosis and treatment of patients with rhinosinusitis and nasal polyps.

2-2 Clinical definition

2-2-1 Clinical definition of rhinosinusitis/nasal polyps

2-2-1-1 Bacteria

Rhinosinusitis (including nasal polyps) is defined as:

- inflammation of the nose and the paranasal sinuses characterised by two or more symptoms, one of which should be either nasal blockage/obstruction/congestion or nasal discharge (anterior/posterior nasal drip):
 - ± facial pain/pressure,
 - ± reduction or loss of smell;

and either

- endoscopic signs of:
 - polyps and/or;
 - mucopurulent discharge primarily from middle meatus and/or; oedema/mucosal obstruction primarily in middle meatus,

and/or


- CT changes:
 - mucosal changes within the ostiomeatal complex and/or sinuses.

2-2-2 Severity of the disease

The disease can be divided into MILD, MODERATE and SEVERE based on total severity visual analogue scale (VAS) score (0 - 10 cm):

- MILD = VAS 0-3
- MODERATE = VAS >3-7
- SEVERE = VAS >7-10

To evaluate the total severity, the patient is asked to indicate on a VAS the answer to the question:



A VAS > 5 affects patient QOL⁽¹⁴⁾.

2-2-3 Duration of the disease

Acute

< 12 weeks

complete resolution of symptoms.

Chronic

> 12 weeks symptoms

without complete resolution of symptoms.

Chronic rhinosinusitis may also be subject to exacerbations

2-3 Definition for use in epidemiology studies/General Practice

For epidemiological studies the definition is based on symptomatology without ENT examination or radiology.

Acute rhinosinusitis (ARS) is defined as:

sudden onset of two or more symptoms, one of which should be either nasal blockage/obstruction/congestion or nasal discharge (anterior/posterior nasal drip):

- ± facial pain/pressure,
- ± reduction or loss of smell;

for < 12 weeks;

with symptom free intervals if the problem is recurrent;

with validation by telephone or interview.

Questions on allergic symptoms i.e. sneezing, watery rhinorrhea, nasal itching and itchy watery eyes should be included.

Acute rhinosinusitis can occur once or more than once in a defined time period. This is usually expressed as episodes/year but there must be complete resolution of symptoms between episodes for it to constitute genuine recurrent acute rhinosinusitis.

Common cold/ acute viral rhinosinusitis is defined as:
duration of symptoms for less than 10 days.

Acute non-viral rhinosinusitis is defined as:
increase of symptoms after 5 days or persistent symptoms after 10 days with less than 12 weeks duration.

Chronic rhinosinusitis with or without nasal polyps is defined as:
presence of two or more symptoms one of which should be either nasal blockage/obstruction/congestion or nasal discharge (anterior/posterior nasal drip):

- ± facial pain/pressure;
- ± reduction or loss of smell;

for > 12 weeks;

with validation by telephone or interview.

Questions on allergic symptoms i.e. sneezing, watery rhinorrhea, nasal itching and itchy watery eyes should be included.

2-4 Definition for research

For research purposes acute rhinosinusitis is defined as above. Bacteriology (antral tap, middle meatal tap) and/or radiology (X-ray, CT) are advised, but not obligatory.

For research purposes chronic rhinosinusitis (CRS) is defined as above. CRS is the major finding and nasal polyposis (NP) is considered a subgroup of this entity. For the purpose of a study, the differentiation between CRS and NP must be based on out-patient endoscopy. The research definition is based on the presence of polyps and prior surgery.

2-4-1 Definition of chronic rhinosinusitis when no earlier sinus surgery has been performed

Chronic rhinosinusitis with nasal polyposis:
polyps bilateral, endoscopically visualised in middle meatus

Chronic rhinosinusitis without nasal polyps:
no visible polyps in middle meatus, if necessary following decongestant

This definition accepts that there is a spectrum of disease in CRS which includes polypoid change in the sinuses and/or middle meatus but excludes those with polypoid disease presenting in the nasal cavity to avoid overlap.

2-4-2 Definition of chronic rhinosinusitis when sinus surgery has been performed

Once surgery has altered the anatomy of the lateral wall, the presence of polyps is defined as bilateral pedunculated lesions as opposed to cobblestoned mucosa > 6 months after surgery on endoscopic examination. Any mucosal disease without overt polyps should be regarded as CRS.

2-4-3 Conditions for sub-analysis

The following conditions should be considered for sub-analysis:

1. aspirin sensitivity based on positive oral, bronchial or nasal provocation or an obvious history;
2. asthma/bronchial hyper-reactivity /COPD/ bronchiectasies based on symptoms, respiratory function tests;
3. allergy based on specific serum IgE or SPT's.

2-4-4 Exclusion from general studies

Patients with the following diseases should be excluded from general studies, but may be the subject of a specific study on chronic rhinosinusitis and/or nasal polyposis:

1. cystic fibrosis based on positive sweat test or DNA alleles;
2. gross immunodeficiency (congenital or acquired);
3. congenital mucociliary problems eg primary ciliary dyskinesia (PCD);
4. non-invasive fungal balls and invasive fungal disease;
5. systemic vasculitis and granulomatous diseases;
6. cocaine abuse;
7. neoplasia.

3. Chronic rhinosinusitis with or without nasal polyps

3-1 Anatomy and (patho)physiology

The nose and paranasal sinuses constitute a collection of air-filled spaces within the anterior skull. The paranasal sinuses communicate with the nasal cavity through small apertures. The nasal cavity and its adjacent paranasal sinuses are lined by pseudostratified columnar ciliated epithelium. This contains goblet cells and nasal glands, producers of nasal secretions that keep the nose moist and form a “tapis roulant” of mucus. Particles and bacteria can be caught in this mucus, rendered harmless by enzymes like lysozyme and lactoferrin, and be transported down towards the oesophagus. Cilia play an important role in mucus transport. All paranasal sinuses are normally cleared by this mucociliary transport, even though transport from large areas of sinuses passes through small openings towards the nasal cavity.

A fundamental role in the pathogenesis of rhinosinusitis is played by the ostiomeatal complex, a functional unit that comprises maxillary sinus ostia, anterior ethmoid cells and their ostia, ethmoid infundibulum, hiatus semilunaris and middle meatus. The key element is the maintenance of the ostial patency. Specifically, ostial patency significantly affects mucus composition and secretion; moreover, an open ostium allows mucociliary clearance to easily remove particulate matter and bacteria. Problems occur if the orifice is too small for the amount of mucus, if mucus production is increased, for instance during an upper respiratory tract infection (URTI), or if ciliary function is impaired. Stasis of secretions follows and bacterial export ceases, causing or exacerbating inflammation of the mucosa whilst aeration of the mucosa is decreased, causing even more ciliary dysfunction. This vicious cycle can be difficult to break, and if the condition persists, it can result as chronic rhinosinusitis. In chronic rhinosinusitis the role of ostium occlusion seems to be less pronounced than in ARS.

3-2 Rhinosinusitis

Rhinosinusitis is an inflammatory process involving the mucosa of the nose and one or more sinuses. The mucosa of the nose and sinuses form a continuum and thus more often than not the mucous membranes of the sinus are involved in diseases which are primarily caused by an inflammation of the nasal mucosa. Chronic rhinosinusitis is a multifactorial disease⁽¹⁵⁾. Factors contributing can be mucociliary impairment^(16, 17), (bacterial) infection⁽¹⁸⁾, allergy⁽¹⁹⁾, swelling of the mucosa for another reason, or rarely physical obstructions caused by morphological/anatomical variations in the nasal cavity or paranasal sinuses^(20, 21). A role in the pathogenesis of rhinosinusitis is certainly played by the ostiomeatal complex, a func-


tional unit that comprises maxillary sinus ostia, anterior ethmoid cells and their ostia, ethmoid infundibulum, hiatus semilunaris and middle meatus. The key element is the maintenance of the ostial patency. An in depth discussion on factors contributing to chronic rhinosinusitis and nasal polyps can be found in chapter 4.

3-3 Chronic rhinosinusitis with or without nasal polyps

Chronic rhinosinusitis with or without nasal polyps is often taken together as one disease entity, because it seems impossible to clearly differentiate both entities⁽²²⁻²⁴⁾. Chronic rhinosinusitis with nasal polyps (CRS without NP) is considered a subgroup of chronic rhinosinusitis (CRS) (Figure 3-1).

The question remains as to why “ballooning” of mucosa develops in polyposis patients and not in all rhinosinusitis patients. Nasal polyps have a strong tendency to recur after surgery even when aeration is improved⁽²⁵⁾. This may reflect a distinct property of the mucosa of polyp patients which has yet to be identified. Some studies have tried to divide chronic rhinosinusitis and nasal polyps based on inflammatory markers⁽²⁶⁻³⁰⁾. Although these studies point to a more pronounced eosinophilia and IL-5 expression in nasal polyps than that found in patients with chronic rhinosinusitis, these studies also point to a continuum in which differences might be found at the ends of the spectrum but at the moment no clear cut division can be made.

Figure 3-1. The spectrum of chronic rhinosinusitis and nasal polyps



Nasal polyps appear as grape-like structures in the upper nasal cavity, originating from within the ostiomeatal complex. They consist of loose connective tissue, oedema, inflammatory cells and some glands and capillaries, and are covered with varying types of epithelium, mostly respiratory pseudostratified epithelium with ciliated cells and goblet cells. Eosinophils are the most common inflammatory cells in nasal polyps, but neutrophils, mast cells, plasma cells, lymphocytes and monocytes are also present, as well as fibroblasts. IL-5 is the predominant cytokine in nasal polyposis, reflecting activation and prolonged survival of eosinophils⁽³¹⁾.

The reason why polyps develop in some patients and not in others remains unknown. There is a definite relationship in patients with 'Samter triad': asthma, NSAID sensitivity and nasal polyps. However, not all patients with NSAID sensitivity have nasal polyps, and vice-versa. In the general population, the prevalence of nasal polyps is 4%⁽³²⁾. In patients with asthma, a prevalence of 7 to 15% has been noted whereas, in NSAID sensitivity, nasal polyps are found in 36 to 60% of

patients^(33, 34). It had long been assumed that allergy predisposed to nasal polyps because the symptoms of watery rhinorrhoea and mucosal swelling are present in both diseases, and eosinophils are abundant. However, epidemiological data provide no evidence for this relationship: polyps are found in 0.5 to 1.5% of patients with positive skin prick tests for common allergens^(34, 35).

4. Epidemiology and predisposing factors

4-1 Introduction

Rhinosinusitis in its many forms, constitutes one of the commonest conditions encountered in medicine and may present to a wide range of clinicians from primary care to accident and emergency, pulmonologists, allergists, otorhinolaryngologists and even intensivists and neurosurgeons when severe complications occur.

The incidence of acute viral rhinosinusitis (common cold) is very high. It has been estimated that adults suffer 2 to 5 colds per year, and school children may suffer 7 to 10 colds per year. The exact incidence is difficult to measure because most patients with common cold do not consult a doctor. Recently a case control study in the Dutch population concluded that an estimated 900000 consultations take place annually for acute respiratory tract infection. Rhinovirus (24%) and Influenzae (11%) were the most common agents isolated.⁽³⁶⁾ More reliable data are available on ARS. As mentioned earlier acute non-viral rhinosinusitis (ARS) is defined as an increase of symptoms after 5 days or persistent symptoms after 10 days after a sudden onset of two or more of the symptoms: nasal blockage/congestion, anterior discharge/postnasal drip, facial pain/pressure, and/or reduction/loss of smell. It is estimated that only 0.5% to 2% of viral URTIs are complicated by bacterial infection; however, the exact incidence is unknown given the difficulty distinguishing viral from bacterial infection without invasive sinus-puncture studies. Bacterial culture results in suspected cases of acute community-acquired sinusitis are positive in only 60% of cases⁽³⁷⁾. Signs and symptoms of bacterial infection may be mild and often resolve spontaneously^(38,39).

In spite of the high incidence of ARS and prevalence and significant morbidity of chronic rhinosinusitis (CRS), with and without nasal polyps, there is only limited accurate data on the epidemiology of these conditions. This observation mainly relates to the lack of a uniformly accepted definition for CRS. In addition, patient selection criteria greatly differ between epidemiologic studies complicating comparison of studies. When interpreting epidemiologic data, one should be aware of a significant selection bias of the different studies presented below. The purpose of this section of the EP3OS document is to give an updated overview of the currently available epidemiologic data on ARS and CRS with and without nasal polyps, and illustrate the factors which are believed to predispose to their development.

4-2 Acute bacterial rhinosinusitis.

When describing the incidence of acute bacterial rhinosinusitis there has been a lot of debate about the actual definition of the

condition. For example in the Cochrane Review on antibiotics for ARS, studies were included if sinusitis was proven by a consistent clinical history, and radiographic or aspiration evidence of ARS⁽⁴⁰⁾. However, most guidelines on the diagnosis of acute bacterial rhinosinusitis base the diagnosis on symptoms and clinical examination. However, if the diagnosis is based on clinical examination alone, the rate of false positive results is high. In patients with a clinical diagnosis of ARS, less than half have significant abnormalities at X-ray examination⁽⁴¹⁾. Based on sinus puncture/aspiration (considered the most accurate), 49-83% of symptomatic patients had ARS⁽⁴²⁾. Compared with puncture/aspiration, radiography offered moderate ability to diagnose sinusitis. Using sinus opacity or fluid as the criterion for sinusitis, radiography had sensitivity of 0.73 and specificity of 0.80⁽⁴²⁾.

An average of 8.4 % of the Dutch population reported at least one episode of ARS per year in 1999⁽⁴³⁾. The incidence of visits to the general practitioner for acute sinusitis in the Netherlands in 2000 was 20 per 1000 men and 33.8 per 1000 women⁽⁴⁴⁾. According to National Ambulatory Medical Care Survey (NAMCS) data in the USA, sinusitis is the fifth most common diagnosis for which an antibiotic is prescribed. Written in 2002, sinusitis accounted for 9% and 21% of all paediatric and adult antibiotic prescriptions respectively⁽⁴⁾.

4-3 Factors associated with acute rhinosinusitis.

4-3-1 Pathogens

Superinfection by bacteria on mucosa damaged by viral infection (common cold) is the most important cause of ARS. The most common bacterial species isolated from the maxillary sinuses of patients with ARS are *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Moraxella catarrhalis*, the latter being more common in children^(45,46). Other streptococcal species, anaerobic bacteria and *Staphylococcus aureus* cause a small percentage of cases. Resistance patterns of the predominant pathogens vary considerably^(47,48). The prevalence and degree of antibacterial resistance in common respiratory pathogens are increasing worldwide. In France, increases in resistance have been observed during the last twenty years in the same geographical area for *H. influenzae* and *S. pneumoniae*⁽⁴⁹⁾. The association between inappropriate antibiotic consumption and the prevalence of resistance is widely assumed based on in vitro experience⁽⁵⁰⁾. Pathogens may also influence the severity of symptomatology⁽⁵¹⁾.

4-3-2 Ciliary impairment

Normal mucociliary flow is a significant non-specific defence mechanism in the prevention of ARS. Viral rhinosinusitis

